



Contribution ID: 16

Type: not specified

Studying Short-Range Nuclear Forces short range correlations through ρ_0 photoproduction at Jefferson Lab

Short Range Correlations (SRCs) are a phenomenon found in all nuclei where two nucleons form a strongly interacting, close-proximity pair in the nucleus, leading to a large relative momentum between the nucleons. Electron-scattering experiments, many of them conducted at Jefferson Lab, have determined that the prevalence of SRCs increase with nuclear size, and furthermore that most SRCs form between a neutron and a proton, a property called 'np-dominance.' Since these observations have largely come from the same type of experiment, it is possible that they are biased by reaction-specific effects, for example, final state interactions. To test the validity of previous observations, an experiment was conducted in Hall D in Fall 2021 using a photon beam on deuterium, helium, and carbon targets to probe SRCs through photoproduction reactions, a radically different approach to test the validity of many previous SRC observations. A preliminary look at the data shows the preliminary results show clear signatures of SRCs, marking the first time that SRCs have been isolated in photoproduction. Using the ρ_0 reaction channel, I plan to compare the rate of proton-proton SRC pairs to that of one proton SRC pairs to test for np-dominance. To assess relative abundances of SRC pairs between heavier targets, I will compare the rates of SRC events from ρ_0 photoproduction in carbon and helium to deuterium. I will also show how these results compare to Monte Carlo simulations.

Primary author: SHARP, Phoebe (George Washington University)

Presenter: SHARP, Phoebe (George Washington University)

Session Classification: Research Talks

Track Classification: Student and Postdoc Talks: Student and Postdoc Talks