



Contribution ID: 333

Type: **Oral Presentation**

Complex potential at $T>0$ from fine lattices

Tuesday, August 9, 2022 5:30 PM (20 minutes)

We study the Wilson line correlation function in Coulomb gauge on $96^3 \times N_t$ lattices in 2+1 flavor QCD with physical strange quark and light quark masses corresponding to pion mass of 310 MeV, with the aim to determine the complex potential at non-zero temperature. In our calculation we use HISQ action in fixed scale approach with lattice spacing $1/a=7.1$ GeV and $N_t=56,36,32,28,24,20$, corresponding to temperatures $T=127, 197, 221, 253, 296, 354$ MeV, respectively. To reduce the noise in the correlator calculations we apply gradient flow. From the analysis of the Wilson line correlation function we conclude that the corresponding spectral function is well described by a dominant peak. The peak position corresponds to the real part of the potential, while the effective width of the peak gives the imaginary part of the potential. We find that the real part of the potential is temperature independent and shows no sign of screening, while the imaginary part shows a strong temperature dependence.

Primary author: PARKAR, Gaurang (University Of Stavanger)

Co-authors: ROTHKOPF, Alexander (University of Stavanger); BAZAVOV, Alexei (Michigan State University); HOYING, Daniel (UConn/BNL); WEBER, Johannes Heinrich (Humboldt-Universität zu Berlin); KACZMAREK, Olaf (University of Bielefeld); PETRECZKY, Peter (BNL); LARSEN, Rasmus (University of Stavanger); MUKHERJEE, Swagato (BNL)

Presenter: PARKAR, Gaurang (University Of Stavanger)

Session Classification: Non-zero Temperature

Track Classification: QCD at Non-zero Temperature