



Contribution ID: 131

Type: **Oral Presentation**

The momentum sum rule via the Feynman Hellmann theorem

Friday 12 August 2022 17:00 (20 minutes)

Determining the quark and gluon contributions to the momentum of a hadron is a difficult and computationally expensive problem. This difficulty mainly arises from the calculation of the gluon matrix element which involves a quark-line disconnected gluon operator, which suffers from noisy ultra-violet fluctuations. Furthermore, a complete calculation also requires a determination of the non-perturbative renormalisation of this operator. In this work, we performed a quenched QCD study of the fully-renormalised quark and gluon contributions to the pion and nucleon momenta via an adaption of the Feynman-Hellmann technique. We find the momentum sum rules are satisfied within our uncertainties for both the pion and nucleon for 3 different values of the quark masses. We also discuss some recent progress on extending this procedure to a dynamical simulations.

Author: ZANOTTI, James (University of Adelaide)

Co-authors: YOUNG, Ross (The University of Adelaide); HORSLEY, Roger (University of Edinburgh); Dr RAKOW, Paul (University of Liverpool); Mr HOWSON, Tomas (University of Adelaide); Dr NAKAMURA, Yoshifumi (RIKEN); Prof. PERLT, Holger (University of Leipzig); Dr STUEBEN, Hinnerk (University of Hamburg); SCHIERHOLZ, Gerrit (DESY)

Presenter: ZANOTTI, James (University of Adelaide)

Session Classification: Hadron Structure

Track Classification: Hadron Structure